

Spectral Conditions for Existence and Uniqueness of Recursive Utilities

Jaroslav Borovička^a and John Stachurski^b

^a Department of Economics, New York University

^b Research School of Economics, Australian National University

October 16, 2017

ABSTRACT. We study existence, uniqueness and computability of solutions for a class of discrete time recursive utilities models. By combining two streams of the recent literature on recursive preferences—one that analyzes principal eigenvalues of valuation operators and another that exploits the theory of monotone concave operators—we obtain conditions that are both necessary and sufficient for existence and uniqueness of solutions. We also show that the natural iterative algorithm is convergent if and only if a solution exists. Consumption processes are allowed to be nonstationary.

JEL Classifications: D81, G11

Keywords: Recursive preferences, existence, uniqueness

1. INTRODUCTION

Recursive preference models such as those discussed in [Koopmans \(1960\)](#), [Epstein and Zin \(1989\)](#) and [Weil \(1990\)](#) play an important role in macroeconomic and financial modeling. For example, the long-run risk models analyzed in [Bansal and Yaron \(2004\)](#), [Hansen et al. \(2008\)](#), [Bansal et al. \(2012\)](#) and [Schorfheide et al. \(2017\)](#) have employed such preferences in discrete time infinite horizon settings with a variety of consumption path specifications to help resolve long-standing empirical puzzles identified in the literature.

In recursive utilities models, the lifetime value of a consumption stream from a given point in time is typically expressed as the solution to a nonlinear forward-looking equation. While this approach is convenient and intuitive, it can be also be

The authors thank Ippei Fujiwara, Jinill Kim, Daisuke Oyama, and Guanlong Ren for many helpful comments. Financial support from ARC grant FT160100423 is gratefully acknowledged.

vacuous, in the sense that no finite solution to the forward looking recursion exists. Moreover, even when a solution does exist, this solution lacks predictive content unless some form of uniqueness can also be established. In general, identifying restrictions that imply existence and uniqueness of a solution for an empirically relevant class of consumption streams is a nontrivial problem.

The best case is where (a) researchers have at hand results implying both existence and uniqueness of solutions, (b) these results are *tight*, and (c) the conditions upon which the results are based are testable in practice. Here tightness means that the underlying conditions are weak enough to accommodate the models and parameterizations that quantitative researchers actually use. Ideally, the conditions will be necessary as well as sufficient, and hence able to exactly characterize the set of well-behaved parameterizations associated with a given class of models.¹

While results on existence and uniqueness of solutions to recursive preference models can be found in the literature, they fall short of the ideal scenario depicted in (a)–(c) above. In particular, there is a gap between the theory they contain and the parameterizations of recursive utility and specifications of consumption paths actually encountered in empirical work at the current research frontier. This gap manifests itself in conditions that are either excessively tight or difficult to test in practice.

The aim of the present paper is to close this gap by obtaining existence and uniqueness results that are as tight as possible in a range of empirically plausible settings, while restricting attention to practical conditions that can be tested in applied work. More specifically, we provide conditions for existence and uniqueness of solutions to the class of preferences studied in [Epstein and Zin \(1989\)](#), while admitting consumption paths that follow a general multiplicative functional specification (see, e.g., [Hansen and Scheinkman, 2009, 2012](#)). These conditions are both necessary and sufficient, and hence as tight as possible in the setting we consider.

To give more detail on that setting, let preferences be defined recursively by

$$(1) \quad V_t = \left[(1 - \beta)C_t^{1-1/\psi} + \beta \{ \mathcal{R}_t(V_{t+1}) \}^{1-1/\psi} \right]^{1/(1-1/\psi)},$$

¹Points (b) and (c) are related: tighter results require weaker conditions, which can be difficult to test in applications. For example, a tight characterization of the set of models for which a solution exists is: the set of models for which a solution exists. This empty truism reminds us that point (c) provides the discipline. Conditions should be testable as well as weak.

where $\{C_t\}$ is a consumption path, V_t is the utility value of the path extending on from time t , and \mathcal{R}_t is the Kreps–Porteus certainty equivalent operator

$$(2) \quad \mathcal{R}_t(V_{t+1}) := (\mathbb{E}_t V_{t+1}^{1-\gamma})^{1/(1-\gamma)}.$$

The parameter $\beta \in (0, 1)$ is a time discount factor, while γ governs risk aversion and ψ is the elasticity of intertemporal substitution (EIS). To ensure that (1) and (2) are well defined, γ and ψ are required to be distinct from 1.

We assume that consumption growth can be expressed as

$$(3) \quad \ln(C_{t+1}/C_t) = \kappa(X_{t+1}, Y_{t+1}, X_t),$$

where κ is a continuous real-valued function, $\{X_t\}$ is a time homogeneous Markov process and $\{Y_t\}$ is an IID innovation process. The persistent component $\{X_t\}$ is required to be compact-valued, while the innovation $\{Y_t\}$ is allowed to be unbounded.

We seek a solution to normalized utility V_t/C_t expressed as a function of the state X_t . Our conditions for existence and uniqueness feature two quantities. The first is the composite parameter

$$(4) \quad \theta := \frac{1 - \gamma}{1 - 1/\psi}.$$

The other is $r(K)$, the spectral radius of a valuation operator K determined by the primitives in (1)–(3) and defined below. We show that when $\theta < 0$, a solution exists if and only if $r(K) > 1$. The same condition is also equivalent to uniqueness of the solution. When $\theta > 0$, the required inequality reverses, and a unique solution exists if and only if $r(K) < 1$. If this condition fails, then no solution exists.

We also show that the above conditions characterizing existence and uniqueness of solutions also imply global convergence of the most natural successive approximations method. Thus, our results also supply a robust, globally convergent method for computing solutions. In fact we show that, if the successive approximations converges, then a unique solution exists and this solution is equal to the limit of the successive approximations. In all other situations, successive approximation will diverge. Thus, if computing the solution to the model at a given set of parameters is the primary objective, then convergence of the iterative method itself justifies the claim that the limit is a solution, and no other solution exists in the candidate space that we consider.

Our work builds on a growing literature on solutions to recursive preference models. The groundwork for Epstein–Zin preferences over infinite-horizon consumption streams was provided by [Epstein and Zin \(1989\)](#), who in turn built on the finite-horizon framework of temporal lotteries found in [Kreps and Porteus \(1978\)](#). [Epstein and Zin \(1989\)](#) obtained sufficient conditions for existence across a broad set of parameters, while allowing geometric consumption growth and eschewing a Markov assumption. These findings were strengthened by [Marinacci and Montrucchio \(2010\)](#), who provided sufficient conditions for both existence and uniqueness of solutions, as well as convergence of successive approximations. Their results were obtained via an innovative fixed point approach that exploits concavity and monotonicity properties possessed by Epstein–Zin preferences with empirically plausible parameterizations.

One issue with the conditions of [Epstein and Zin \(1989\)](#) and [Marinacci and Montrucchio \(2010\)](#) is that they require, at least asymptotically, a finite bound b on consumption growth $\ln(C_{t+1}/C_t)$ that holds with probability one. This fails for many standard consumption processes. One example is [Bansal and Yaron \(2004\)](#), where

$$(5) \quad \ln(C_{t+1}/C_t) = \mu + z_t + \sigma_t \eta_{t+1}.$$

Here $\{z_t\}$ and $\{\sigma_t\}$ are stationary processes and $\{\eta_{t+1}\}$ is IID and $N(0, 1)$. Evidently consumption growth is unbounded above.

In fact the problem is not so much that the bounded growth condition fails, since the shocks in (5) can be truncated at suitably large values and b can then be chosen as the maximal value taken by the right-hand side of (5). Rather, the issue is that the resulting restrictions on preference parameters, which are used to ensure finiteness of the solution to the recursive utility model, are excessively conservative. The intuition behind this is that the probability one bound b considers only the worst case in terms of pushing utility to infinity. Much sharper results can be obtained by considering what happens “on average.” After all, recursive utility specifications, while nonlinear, are still defined using expectations. This means that the whole distribution matters, not just the right-hand tail.

As stated above, the conditions presented in this paper are based around the spectral radius of the operator K , which captures the distributions of the persistent components and innovations in consumption specifications such as (5). Since the spectral radius depends on the full distributions and involves averaging by integration, it leads to conditions that are much sharper—in fact as sharp as possible for the setting we consider, being both necessary and sufficient. By applying

our results to two empirical set ups from the recent literature ([Bansal and Yaron \(2004\)](#), [Schorfheide et al. \(2017\)](#)), we demonstrate that the gap between the necessary and sufficient conditions developed here and the sufficient conditions arising from probability one bounds is both large and significant for modern quantitative applications.

We should add two qualifications to the preceding discussion. First, [Marinacci and Montrucchio \(2010\)](#) treat a much larger range of recursive utility specifications than we consider here, as well as admitting non-Markovian state processes. Thus, while our results are sharper for the problems we consider, their study is far more comprehensive. Second, our condition is inherently computational, since the spectral radius cannot be derived analytically apart from very special cases. The computational—as opposed to analytical—nature of our condition facilitates tight results.

In addition to ourselves, a number of other researchers have sought to extend the work of [Marinacci and Montrucchio \(2010\)](#). In the realm of recursive utility specifications with Thompson aggregators, the idea of exploiting monotonicity and concavity of the Koopmans operator has been adapted and extended by [Balbus \(2016\)](#), [Becker and Rincón-Zapatero \(2017\)](#) and [Bloise and Vailakis \(2017\)](#). While these contributions do not resolve the issues associated with using probability one bounds on consumption growth discussed above, they carefully elucidate the links between the monotonicity and concavity properties of certain aggregators and fixed point results in partially ordered vector spaces. Such results also lie at the heart of this paper, and we have benefited extensively from their ideas.²

There is one more closely related paper upon which we draw extensively: the study of Epstein–Zin utility models with fully unbounded consumption growth specifications in [Hansen and Scheinkman \(2012\)](#). Their approach is to draw a connection between the solution to the Epstein–Zin utility recursion and the Perron–Frobenius eigenvalue problem associated with a linear operator, denoted in their paper by \mathbb{T} , that is proportional to the operator K discussed above. Consumption

²Prior to [Marinacci and Montrucchio \(2010\)](#), important contributions to the literature on existence and uniqueness of solutions to recursively defined utility specifications (as well as the closely related problem of optimality of dynamic programs with general aggregators), were made by [Koopmans \(1960\)](#), [Lucas and Stokey \(1984\)](#), [Becker et al. \(1989\)](#), [Streufert \(1990\)](#), [Boyd \(1990\)](#), [Ozaki and Streufert \(1996\)](#), [Le Van and Vailakis \(2005\)](#) and [Rincón-Zapatero and Rodríguez-Palmero \(2007\)](#). We also note the important contributions of earlier authors who used monotonicity and concavity to obtain fixed points of forward looking recursive models. See, for example, [Coleman \(1991\)](#), [Datta et al. \(2002\)](#) and [Mirman et al. \(2008\)](#).

growth obeys (3) and, unlike this paper, $\{X_t\}$ is not required to be compact-valued. In this very general setting they show that a solution exists when a joint restriction holds on the spectral radius of \mathbb{T} and the preference parameters, in addition to as certain auxiliary restrictions. They also obtain a uniqueness result for the case $\theta \geq 1$.³

The advantage of the conditions in [Hansen and Scheinkman \(2012\)](#) is the lack of a compactness restriction on X_t . The advantages of our approach are as follows: First, we obtain uniqueness of the solution for all θ , not just for $\theta \geq 1$. (One reason this matters is that empirical studies typically find that $\theta < 1$.) Second, we obtain conditions that are necessary as well as sufficient, both for existence and for uniqueness. Third, we obtain a globally convergent method of computation, and show that it converges if and only if a solution exists. Fourth, the auxiliary conditions in [Hansen and Scheinkman \(2012\)](#), which generalize our compactness assumption, involve testing integrability restrictions on the eigenfunctions of the operator \mathbb{T} . In general these kinds of conditions are difficult to test, unless, of course, one truncates the state space—which is exactly what we do here.⁴

While compactness must be imposed on persistent components of consumption growth in order to use our conditions, these components are bounded in probability, and hence one can always choose a compactification of the state space such that the overall impact on the stochastic process for consumption is arbitrarily small. Computational results in section 3.3 show that, for the applications we consider, the spectral radius calculation results are stable with respect to truncation of the innovation terms whenever the truncation points are sufficiently large.

The other connection between our paper and that of [Hansen and Scheinkman \(2012\)](#) is both sets of results rely on information contained in the dominant eigenvalue and corresponding eigenfunction of the valuation operator. Indeed, the principal eigenpairs of valuation operators associated with future cash and utility payoffs have increasingly been used to understand long run risks and long run values in macroeconomic and financial applications (see, e.g., [Alvarez and Jermann \(2005\)](#); [Hansen and Scheinkman \(2009\)](#); [Qin and Linetsky \(2017\)](#)). In this paper we connect the principal eigenpairs of the valuation operator associated with Epstein–Zin preferences to the theory of monotone concave operators. In this way we link

³See proposition 6 of [Hansen and Scheinkman \(2012\)](#). Note that the symbol α in their study corresponds to θ here.

⁴Another recent paper that works in a noncompact state setting is [Christensen \(2017\)](#). There the focus is on robust decision makers, who can also be viewed as utility maximizers with risk-sensitive preferences.

two active strands of research on present values associated with cash and utility flows.⁵

The paper is structured as follows: Section 2 states our main results. Section 3 discusses applications. Section 4 begins the process of providing proofs by developing a set of fixed point results in an abstract setting. Section 5 completes the proofs by connecting these abstract results to the Epstein–Zin recursive utility setting.

2. MAIN RESULTS

To maintain focus on results and applications, we first state all findings on existence, uniqueness and convergence of successive approximations for Epstein–Zin recursive preference models.

2.1. Definitions and Assumptions. Let consumption $\{C_t\}$ be as defined in (3), where the state process $\{X_t\}$ is time homogeneous and Markovian, taking values in some metric space \mathbb{X} . The stochastic kernel for $\{X_t\}$ will be denoted by Q . In particular,

$$Q(x, B) = \mathbb{P}\{X_{t+1} \in B \mid X_t = x\}$$

for all $x \in \mathbb{X}$ and all Borel sets $B \subset \mathbb{X}$. The innovation process $\{Y_t\}$ is IID, independent of $\{X_t\}$, and takes values in some topological space \mathbb{Y} . The common distribution of each Y_t is a Borel probability measure on \mathbb{Y} denoted by ν . Throughout we impose the following restrictions.

Assumption 2.1. \mathbb{X} is compact and the growth function κ is continuous.

Let \mathcal{C} be all continuous real-valued functions on \mathbb{X} and let $\|\cdot\|$ be the supremum norm on \mathcal{C} . Let \mathcal{C}_+ be all nonnegative functions in \mathcal{C} . A linear operator K from \mathcal{C} to itself is called *strongly positive* if Kg is strictly positive everywhere on \mathbb{X} whenever $g \in \mathcal{C}_+$ and g is nonzero. K is called *compact* if the image under K of the unit ball in \mathcal{C} is relatively compact. The *operator norm* and *spectral radius* of K are defined by

$$(6) \quad \|K\| := \sup\{\|Kg\| : g \in \mathcal{C}, \|g\| \leq 1\} \quad \text{and} \quad r(K) := \lim_{n \rightarrow \infty} \|K^n\|^{1/n}$$

respectively.⁶

⁵We also connect analysis of principle eigenvalues to a version of Banach’s contraction mapping theorem, which is needed for the case $\theta \in (0, 1)$.

⁶While the spectral radius is sometimes defined as the supremum of the modulus of elements of the spectrum of K , these two definitions are equivalent for linear operators on \mathcal{C} . See, for example, [Kolmogorov and Fomin \(1975\)](#).

2.2. The Fixed Point Problem. Our interest centers on existence, uniqueness and computability of V_t in (1)–(2). We convert this into a fixed point problem following the steps in Hansen and Scheinkman (2012). Minor manipulations to (1) yield

$$\left(\frac{V_t}{C_t}\right)^{1-1/\psi} = \zeta + \beta \left\{ \mathcal{R}_t \left(\frac{V_{t+1} C_{t+1}}{C_{t+1} C_t} \right) \right\}^{1-1/\psi},$$

where $\zeta := 1 - \beta$. (The results below are valid for any positive constant ζ .) With $\theta := (1 - \gamma)/(1 - 1/\psi)$ and $W_t := (V_t/C_t)^{1-1/\psi}$, we can rewrite this as

$$(7) \quad W_t = \zeta + \beta \left\{ \mathbb{E}_t W_{t+1}^\theta \exp[(1 - \gamma)\kappa(X_{t+1}, Y_{t+1}, X_t)] \right\}^{1/\theta}.$$

We seek a Markov solution $W_t = w(X_t)$ for some $w: \mathbb{X} \rightarrow \mathbb{R}$, and hence solutions to the functional equation

$$(8) \quad w(x) = \zeta + \beta \left\{ \int w(x')^\theta \int \exp[(1 - \gamma)\kappa(x', y', x)] \nu(dy') Q(x, dx') \right\}^{1/\theta}.$$

While $w \equiv +\infty$ can be considered as solution to (8), we focus on solutions in \mathcal{C}_+ .

Let K be the operator defined on \mathcal{C} by

$$(9) \quad Kg(x) = \beta^\theta \int g(x') \int \exp[(1 - \gamma)\kappa(x', y', x)] \nu(dy') Q(x, dx')$$

and let T be defined on \mathcal{C}_+ by

$$(10) \quad Tw(x) = \zeta + [Kw^\theta(x)]^{1/\theta}.$$

A function $w \in \mathcal{C}_+$ solves (8) if and only if it is a fixed point of T in \mathcal{C}_+ .

Since $\zeta > 0$, T maps nonnegative functions into strictly positive functions.⁷ Hence any fixed point of T must lie in the interior of \mathcal{C}_+ .

Returning to (9), evidently $g \mapsto Kg$ is linear. We add the following restrictions:

Assumption 2.2. K is a strongly positive compact operator from \mathcal{C} to itself.

Assumption 2.2 holds in all applications we consider.

Example 2.1. Suppose that $\mathbb{X} \subset \mathbb{R}^n$ and $Q(x, dx') = q(x, x')dx'$ for some continuous positive density kernel $q: \mathbb{X} \times \mathbb{X} \rightarrow \mathbb{R}$. Then K is an integral operator $Kg(x) = \int g(x')k(x, x')dx'$ with kernel

$$k(x, x') := \beta^\theta \int \exp[(1 - \gamma)\kappa(x', y', x)] \nu(dy') q(x, x').$$

⁷Some ambiguity in (10) arises when $\theta < 0$ and $w(x) = 0$ for some x . The general rule we apply for exponent $\alpha < 0$ is $0^\alpha = \infty$ and $\infty^\alpha = 0$. Thus, when w is not in $\mathring{\mathcal{C}}_+$, we admit $Kw^\theta(x) = \infty$ at some x , and at these points $Tw(x) = \zeta$. When $Kw^\theta(x) < \infty$ we have $Tw(x) > \zeta$.

Evidently $g \geq 0$ implies $Kg \gg 0$. Joint continuity of k combined with compactness of \mathbb{X} implies compactness of K as a linear operator on \mathcal{C} .⁸ Hence assumption 2.2 is valid.

Example 2.2. Consider the setting of example 2.1, except that \mathbb{X} is finite, endowed with the discrete topology, and q is a density with respect to the counting measure instead of Lebesgue measure (i.e., q is a stochastic matrix). The conditions of assumption 2.2 are again satisfied.

2.3. Results for Recursive Utility. We begin with the case $\theta < 0$, which corresponds to most empirically relevant parameterizations. The proof of the following result uses a principal eigenfunction theorem combined with the theory of monotone concave operators. See section 5 for details.

Theorem 2.1. *If $\theta < 0$, then the following statements are equivalent:*

- (a) $r(K) > 1$.
- (b) T has a fixed point in \mathcal{C}_+ .
- (c) There exists a $w \in \mathcal{C}_+$ such that $\{T^n w\}_{n \geq 1}$ is convergent in \mathcal{C}_+ .
- (d) T has a unique fixed point in \mathcal{C}_+ .
- (e) T is globally asymptotically stable on \mathcal{C}_+ .

We turn next to the case $\theta > 0$, where the condition on $r(K)$ reverses.

Theorem 2.2. *If $\theta > 0$, then the following statements are equivalent:*

- (a) $r(K) < 1$.
- (b) T has a fixed point in \mathcal{C}_+ .
- (c) There exists a $w \in \mathcal{C}_+$ such that $\{T^n w\}_{n \geq 1}$ is convergent in \mathcal{C}_+ .
- (d) T has a unique fixed point in \mathcal{C}_+ .
- (e) T is globally asymptotically stable on \mathcal{C}_+ .

Unlike theorem 2.1, the proof of theorem 2.2 has to be broken down into two cases: $0 < \theta < 1$ and $1 \leq \theta$. In the first case the conjugate operator A defined in (22) has the property that A^n is a contraction for some n , but A is not a concave operator. In the case $\theta \geq 1$, the operator A is not generally contracting, but it is a concave operator. See section 5.

⁸See, for example, [Kolmogorov and Fomin \(1975\)](#), §24.

3. APPLICATIONS

Let us apply the results presented above to some specifications from recent empirical studies and test whether or not the conditions of theorems 2.1–2.2 are satisfied. For comparison, we also investigate whether or not the conditions of the tests from the existing literature based on probability one bounds hold. Code that replicates all computations can be found at

https://github.com/jstac/recursive_utility_code.

When checking the conditions of theorems 2.1–2.2, we need to compute the spectral radius of K under each parameterization. We do so by applying a theorem of V. Ya. Stet'senko, reproduced as theorem 9.1 of [Krasnosel'skii et al. \(2012\)](#), which states that if K is linear operator that leaves a solid normal order cone invariant, then

$$(11) \quad r(K) = \lim_{n \rightarrow \infty} \|K^n g\|^{1/n}$$

for any g that lies in the interior of that cone.⁹ Since the valuation operator K defined in (9) maps \mathcal{C}_+ to itself and the latter is both solid and normal under the supremum norm, the formula (11) can be applied by iterating with K starting from any strictly positive continuous function g on \mathbb{X} .

3.1. Long-Run Risk. Suppose first that consumption growth obeys the [Bansal and Yaron \(2004\)](#) long-run risk specification

$$\begin{aligned} \ln(C_{t+1}/C_t) &= \mu + z_t + \sigma_t \eta_{c,t+1}, \\ z_{t+1} &= \rho z_t + \phi_z \sigma_t \eta_{z,t+1}, \\ \sigma_{t+1}^2 &= v \sigma_t^2 + d + \phi_\sigma \eta_{\sigma,t+1}. \end{aligned}$$

Here $\{\eta_{i,t}\}$ are IID and standard normal for $i \in \{c, z, \sigma\}$. The state for consumption can be represented as the Markov process $X_t = (z_t, \sigma_t)$.

Consider the probability one test for existence and uniqueness of solutions based on the conditions in [Epstein and Zin \(1989\)](#) and [Marinacci and Montrucchio \(2010\)](#)

⁹This is significantly easier to calculate than the limit of $\|K^n\|^{1/n}$ given in the definition of $r(K)$, since the latter involves an operator norm. Another alternative is to discretize the state process $\{X_t\}$, reducing K to a matrix, and then compute the spectral radius of the matrix by extracting the eigenvalues using standard algorithms. We also implemented this method and obtained similar results, albeit with some degree of sensitivity to the means of discretization. Discretization based methods also suffer more from increasing dimensionality.

outlined in the introduction. Theorem 3 of [Marinacci and Montrucchio \(2010\)](#) shows that a unique solution to the recursive utility problem exists whenever

$$(12) \quad s(b, \beta, \psi) < 1, \quad \text{where} \quad s(b, \beta, \psi) := \exp(b)\beta^{1/(1-1/\psi)}$$

and b is an almost sure (i.e., probability one) upper bound on $\ln(C_{t+1}/C_t)$. Since consumption growth is in fact unbounded in this model, we bound it by truncating the stationary distribution of consumption growth at the 95th percentile. The upper bound b of log consumption growth is then computed and the expression in (12) is evaluated.¹⁰

The other test we consider is the spectral radius based test in theorems 2.1–2.2. While consumption growth need not be bounded for this test, the state space must be compact before theorems 2.1–2.2 can be applied. In principle this requires truncating the innovations in the state process. In practice truncation is inherent in all computation, since the set of floating point numbers on any machine with finite memory is bounded. (On a 64 bit machine these bounds are very large and never bind.) The integrals in (9) are computed by either Monte Carlo or Gaussian quadrature, with both methods giving similar results. Where necessary, functions are fitted using continuous piecewise linear interpolation.

In [Bansal and Yaron \(2004\)](#), the preference parameters are set to $\gamma = 10.0$, $\beta = 0.998$ and $\psi = 1.5$. Here and in all subsequent cases $\zeta = 1 - \beta$. The parameters in the consumption process are $\mu = 0.0015$, $\rho = 0.979$, $\phi_z = 0.044$, $v = 0.987$, $d = 7.9092e-7$ and $\phi_\sigma = 2.3e-6$. Evaluating the left-hand side of (12) at these parameters using the methodology described above yields 1.0254. Thus, the Bansal–Yaron model fails the probability one-based sufficient condition. Nonetheless, $\theta = -27$ and the spectral radius $r(K)$ evaluates to 1.068. Hence, by theorem 2.1, a unique solution exists. This tells us (i) that the solution to the Bansal–Yaron specification is well-defined, unique and can be computed from any initial condition in \mathcal{C}_+ by successive approximation, and (ii) that the probability one condition is too strict to effectively treat this model.

Figure 1a further illustrates the same point by showing the value of the expression on the left-hand side of (12) at the Bansal–Yaron parameterization and also at neighboring parameterizations obtained by varying ψ and μ . Almost all values

¹⁰While the 95th percentile is a relatively imprecise truncation exceeded 5% of the time in long sample paths, we show below that the condition in (12) is too strict to be constructively applied to the models we consider. Adopting a higher percentile in the truncation only intensifies this problem.



(A) The probability one test value $s(b, \beta, \psi)$ (B) The spectral radius $r(K)$

FIGURE 1. Stability tests for the Bansal–Yaron model

exceed unity, indicating failure of the test, apart from a small measure of parameterizations to the left of the 1.0 contour line. Nonetheless, most of these models are in fact stable, with a unique, globally attracting solution. This is true because $\theta < 0$ in all cases and, as shown in figure 1b, only in the north-east corner do we find parameterizations with $r(K) \leq 1$.

3.2. Schorfheide–Song–Yaron Consumption Dynamics. Next consider the consumption specification adopted in [Schorfheide et al. \(2017\)](#), where

$$\begin{aligned}\ln(C_{t+1}/C_t) &= \mu + z_t + \sigma_{c,t} \eta_{c,t+1}, \\ z_{t+1} &= \rho z_t + \sqrt{1 - \rho^2} \sigma_{z,t} \eta_{z,t+1}, \\ \sigma_{i,t} &= \phi_i \bar{\sigma} \exp(h_{i,t}) \quad \text{with} \quad h_{i,t+1} = \rho_{h_i} h_i + \sigma_{h_i} \eta_{h_i,t+1}, \quad i \in \{c, z\}.\end{aligned}$$

The innovations $\{\eta_{c,t}\}$ and $\{\eta_{h_i,t}\}$ are IID and standard normal for $i \in \{c, z\}$. The state can be represented as $X_t = (\sigma_{c,t}, \sigma_{z,t}, z_t)$.

In [Schorfheide et al. \(2017\)](#), the preference parameters are calibrated to be $\gamma = 8.89$, $\beta = 0.999$ and $\psi = 1.97$, while the parameters in the consumption process are $\mu = 0.0016$, $\rho = 0.987$, $\phi_z = 0.215$, $\bar{\sigma} = 0.0032$, $\phi_c = 1.0$, $\rho_{h_z} = 0.992$, $\sigma_{h_z}^2 = 0.0039$, $\rho_{h_c} = 0.991$, and $\sigma_{h_c}^2 = 0.0096$. Following procedures analogous to those used for the model of [Bansal and Yaron \(2004\)](#), we find that the left-hand side of (12) evaluates to 1.003. Thus, the sufficient condition for existence based on the probability one bound fails to hold. At the same time, $\theta = -16.02$ and, by calculations analogous to those used to compute the spectral radius in the [Bansal and Yaron \(2004\)](#) case, we obtain $r(K) = 1.050$. Hence a unique solution does exist, by theorem 2.1.

Figures 1a–1b illustrate further by repeating the exercise of examining the tests at neighboring parameterizations. The results are similar to those obtained for the Bansal–Yaron parameterization: The probability one based test is too conservative, excluding many parameterizations that do in fact have unique, well-defined solutions. (Here $\theta < 0$ again occurs for all parameter values shown, so $r(K) > 1$ corresponds to the stable case.) A similar outcome is observed in figures 3a–3b, when β is varied instead of μ .

3.3. Truncations Errors. To assess the information loss from compactification, we compute the spectral radius for the [Bansal and Yaron \(2004\)](#) specification at different levels of truncation. More specifically, we replace all of the shocks η_i in section 3.1 with $\hat{\eta}_i$ truncated to lie within the interval $[-\tau, \tau]$. The parameter τ is the variable on the horizontal axis of figure 4. The vertical axis shows the corresponding spectral radius, computed using (11). Small values of τ cause significant errors because they distort the model via the distribution of the innovation terms. As τ increases the computed value of the spectral radius stabilizes. In the preceding computations we used $\tau = 3$.



(A) The probability one test value $s(b, \beta, \psi)$ (B) The spectral radius $r(K)$

FIGURE 2. Stability tests for the Schorfheide–Song–Yaron model

4. PROOFS PART I: GENERAL FIXED POINT RESULTS

We begin with an abstract fixed point problem for an operator A of the form $Ag(x) = \phi(Kg(x))$, where K is a linear operator and ϕ is a scalar function. We obtain a range of fixed point results depending on the properties of ϕ and K . Our motivation is that the recursive preference equation studied in subsequent sections can be mapped to such a fixed point problem after a continuous transformation.



(A) The probability one test value $s(b, \beta, \psi)$ (B) The spectral radius $r(K)$

FIGURE 3. Stability tests for the Schorfheide–Song–Yaron model

In that setting, the operator K corresponds to the valuation operator discussed in the introduction to this paper.

4.1. Preliminaries. Let \mathbb{X} be any compact metric space. As before, let \mathcal{C} be all continuous real-valued functions on \mathbb{X} . Let \leq be the usual pointwise partial order on \mathcal{C} and let $\|\cdot\|$ be the supremum norm. The symbol \ll denotes strict pointwise inequality, so that $f \ll g$ means $f(x) < g(x)$ for all $x \in \mathbb{X}$. The statement $f < g$




FIGURE 4. Spectral radius $r(K)$ as a function of truncation value

means that $f \leq g$ and $f(x) < g(x)$ for some $x \in \mathbb{X}$. Given $f \leq g$ in \mathcal{C} , let

$$[f, g] := \{h \in \mathcal{C} : f \leq h \leq g\}.$$

Note that \mathcal{C} is a Banach lattice with the supremum norm and partial order \leq . In particular, the metric induced by $\|\cdot\|$ on \mathcal{C} is complete, \mathcal{C} is closed under the taking of pairwise suprema and $|f| \leq |g|$ implies $\|f\| \leq \|g\|$ for all f, g in \mathcal{C} .¹¹

The *positive cone* of \mathcal{C} , denoted below by \mathcal{C}_+ , is all $g \in \mathcal{C}$ such that $0 \leq g$. As an order cone in \mathcal{C} , the set \mathcal{C}_+ is solid, normal and reproducing.¹² The interior of \mathcal{C}_+ is denoted $\overset{\circ}{\mathcal{C}}_+$ and contains all $g \in \mathcal{C}$ with $0 \ll g$. An operator A from $\mathcal{C}_0 \subset \mathcal{C}$ into \mathcal{C} is called *increasing* if $Af \leq Ag$ whenever $f, g \in \mathcal{C}_0$ and $f \leq g$. It is called *concave* if, for all $\alpha \in [0, 1]$ and $f, g \in \mathcal{C}_0$,

$$(13) \quad \alpha Af + (1 - \alpha)Ag \leq A\{\alpha f + (1 - \alpha)g\}.$$

Note that, for any linear increasing map L on \mathcal{C} we have $|Lf| \leq L|f|$ for every $f \in \mathcal{C}$.

Given a linear operator L from \mathcal{C} to itself, the *operator norm* and *spectral radius* of L are defined by (6). The definitions of compactness and strong positivity of L are as given in section 2.1.

We will exploit the following fixed point theorem for monotone concave operators:

¹¹See Zaanen (1997) and Aliprantis and Border (2006) for more details.

¹²A cone C in \mathcal{C} is called *solid* if it has nonempty interior, *normal* if there exists a constant N with $\|f\| \leq N\|g\|$ whenever $f, g \in C$ with $f \leq g$, and *reproducing* if every element f of \mathcal{C} can be written as a linear combination of elements of C . See, for example, Du (2006).

Theorem 4.1 (Du–Zhang). *Let A be increasing and concave on \mathcal{C}_+ . If, in addition, there exist functions $f_1 \leq f_2$ in \mathcal{C}_+ with $Af_1 \gg f_1$ and $Af_2 \leq f_2$, then A is globally asymptotically stable on $[f_1, f_2]$.*

Proof. This follows from corollary 2.1.1 of Zhang (2013), given that \mathcal{C}_+ is both normal and solid as an order cone in \mathcal{C} . \square

4.2. Set Up. Let $K: \mathcal{C} \rightarrow \mathcal{C}$ be an increasing linear operator and let $\phi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be continuous. Let Φ be the operator on \mathcal{C}_+ defined by $\Phi g = \phi \circ g$. Let A be the operator on \mathcal{C}_+ defined by $Ag = \Phi Kg$, or, equivalently,

$$(14) \quad (Ag)(x) = \phi(Kg(x)) \quad (x \in \mathbb{X}).$$

Note that $A\mathcal{C}_+ \subset \mathcal{C}_+$. Indeed, given that K is linear and increasing, for each fixed $g \in \mathcal{C}_+$ we have $Kg \in \mathcal{C}_+$. Since ϕ is continuous and nonnegative, ΦKg is also in \mathcal{C}_+ . While A is a nonlinear operator, the nonlinear component enters only as a scalar transformation. Below we consider fixed points of A in \mathcal{C}_+ under a range of auxiliary assumptions. Assumptions placed on K and ϕ in this section are always in force.

4.3. Long Run Contractions. Suppose for the purposes of this section that ϕ is Lipschitz of order ℓ for some $\ell > 0$. That is,

$$(15) \quad |\phi(s) - \phi(t)| \leq \ell |s - t| \text{ for all } s, t \geq 0.$$

First we consider the case where the spectral radius of K is small relative to $1/\ell$.

Proposition 4.1. *If $r(K) < 1/\ell$, then there exists an $n \in \mathbb{N}$ such that A^n is a contraction map on \mathcal{C}_+ . In particular, A is globally asymptotically stable on \mathcal{C}_+ .*

Proof of proposition 4.1. Fix $f, g \in \mathcal{C}_+$. As a first step we claim that

$$(16) \quad |A^n f - A^n g| \leq (\ell K)^n |f - g|$$

holds pointwise on \mathbb{X} for all integers $n \geq 0$. It holds for $n = 0$, since A^0 and K^0 are by definition identity maps. Now suppose that it holds for some $n \geq 0$. We claim it also holds at $n + 1$. Indeed

$$\left| A^{n+1} f - A^{n+1} g \right| = \left| \Phi K A^n f - \Phi K A^n g \right| \leq \ell |K A^n f - K A^n g|,$$

where the inequality is due to (15). Using this bound and the linearity and monotonicity of K leads us to

$$\left| A^{n+1} f - A^{n+1} g \right| \leq \ell |K(A^n f - A^n g)| \leq \ell K |A^n f - A^n g| \leq (\ell K)^{n+1} |f - g|,$$

where the last inequality uses the induction hypothesis combined with the monotonicity of K . We have now confirmed that (16) holds for all $n \geq 0$.

Taking the supremum over (16) yields $\|A^n f - A^n g\| \leq \|(\ell K)^n |f - g|\|$. The definition of the operator norm then gives

$$\|A^n f - A^n g\| \leq \|(\ell K)^n\| \cdot \|f - g\| = \ell^n \|K^n\| \cdot \|f - g\|.$$

From the definition $r(K) = \lim_{n \rightarrow \infty} \|K^n\|^{1/n}$, we have

$$(\ell^n \|K^n\|)^{1/n} \rightarrow \ell r(K) < 1.$$

Hence, for sufficiently large n , the map A^n is a contraction with modulus $\ell^n \|K^n\|$. All claims in proposition 4.1 follow from this property, completeness of $\|\cdot\|$, the fact that \mathcal{C}_+ is closed in \mathcal{C} , and a well-known extension to the Banach contraction mapping theorem (see, e.g., p. 272 of Wagner (1982)). \square

Remark 4.1. Since $r(K) \leq \|K\|$, to obtain the spectral radius condition in proposition 4.1, it suffices to show that $\|K\| < 1/\ell$. In fact, as is clear from the proof of proposition 4.1, A itself becomes a contraction mapping under this condition.

4.4. Fixed Points under Monotonicity and Concavity. If $\phi(0) = 0$, then $0 \in \mathcal{C}_+$ is clearly a fixed point of A . In applications this fixed point is often trivial, and interest centers on positive fixed points. In such settings it turns out that notions of concavity and monotonicity are more helpful than contractivity arguments for establishing fixed point results. We begin with a simple lemma.

Lemma 4.1. *For the operator $A = \Phi K$ defined above, the following statements hold.*

- (a) *If ϕ is increasing on \mathbb{R}_+ , then A is increasing on \mathcal{C}_+ .*
- (b) *If ϕ is concave on \mathbb{R}_+ , then A is concave on \mathcal{C}_+ .*

Proof. Claim (a) is obviously true, given that K is already assumed to be increasing. Regarding claim (b), since K is linear, for any fixed $f, g \in \mathcal{C}_+$, $x \in \mathbb{X}$, and $\alpha \in [0, 1]$, we have

$$\begin{aligned} \alpha \phi(Kf(x)) + (1 - \alpha) \phi(Kg(x)) &\leq \phi(\alpha Kf(x) + (1 - \alpha)Kg(x)) \\ &= \phi[K(\alpha f(x) + (1 - \alpha)g(x))]. \end{aligned}$$

In other words, $\alpha Af(x) + (1 - \alpha)Ag(x) \leq A[\alpha f(x) + (1 - \alpha)g(x)]$ for any $x \in \mathbb{X}$, as was to be shown. \square

Lemma 4.2. *If ϕ is strictly concave, K is strongly positive, and, in addition,*

$$(17) \quad \phi(0) = 0, \quad \lim_{t \downarrow 0} \frac{\phi(t)}{t} \leq 1 \quad \text{and} \quad r(K) \leq 1,$$

then the only fixed point of A in \mathcal{C}_+ is 0.

Proof. Since K is linear we have $K0 = 0$. Hence, with the additional assumption $\phi(0) = 0$, we have $A0 = 0$. Thus, the zero element is a fixed point. It remains to show that no other fixed point exists in \mathcal{C}_+ . In doing so we use the fact that ϕ must be strictly positive everywhere on $(0, \infty)$, since the existence of a positive t with $\phi(t) = 0$ violates our assumption that ϕ is both strictly concave and nonnegative.

Seeking a contradiction, suppose that $g \in \mathcal{C}_+$ is a nonzero fixed point of A . Observe that, since $Ag = g$ and g is nonzero, the fact that K is a strongly positive operator and ϕ is positive on $(0, \infty)$ implies that $g \gg 0$. In particular, the constant $m := \min g$ is strictly positive. Let $\eta := \phi(m)/m$. Note that, by strict concavity of ϕ and the assumption that $\frac{\phi(t)}{t} \rightarrow 1$ as $t \downarrow 0$, we have

$$(18) \quad \eta < 1 \quad \text{and} \quad t \geq m \implies \phi(t) \leq \eta t.$$

Observe that $Kg \gg g$ must hold. To see why, suppose that $Kg(x) \leq g(x)$ for some x . Invoking strict concavity and the limit in (17) again, we have $\phi(t) < t$ for any positive t , and hence $Ag(x) = \phi(Kg(x)) < Kg(x) \leq g(x)$. This contradicts the assumption that g is a fixed point of A . Our claim that $Kg \gg g$ is confirmed.

Next we claim that $A^n g \leq (\eta K)^n g$ for all n . Evidently this holds at $n = 0$, and, assuming it holds at n , we have

$$A^{n+1}g = \Phi K A^n g \leq \eta K A^n g \leq \eta K (\eta K)^n g = (\eta K)^{n+1} g.$$

In the first inequality we used the fact that $Kg \gg g$ and g is a fixed point of A , so that $K A^n g = Kg \gg g \geq m$ when m is as given in (18). In the second inequality we used the induction hypothesis and the monotonicity of K .

We have now shown by induction that $A^n g \leq (\eta K)^n g$ for all $n \in \mathbb{N}$. Hence

$$(19) \quad \|A^n g\| \leq \eta^n \|K^n g\| \leq \eta^n \|K^n\| \|g\|$$

for all n . Since $r(K) \leq 1$ and $\eta < 1$, the definition of $r(K)$ implies existence of an $n \in \mathbb{N}$ such that $\|K^n\|^{1/n} < 1/\eta$, or $\|K^n\| < (1/\eta)^n$. Evaluating (19) at this n gives $\|A^n g\| < \|g\|$, contradicting our assumption that g is a fixed point of A . \square

Below we will make use of the following version of the Krein–Rutman theorem, the value of which for studying recursive preference models was identified and illustrated in [Hansen and Scheinkman \(2009, 2012\)](#).

Lemma 4.3 (Krein–Rutman). *If, in addition to being linear and increasing, K is also strongly positive and compact, then $r(K) > 0$ and $r(K)$ is an eigenvalue of K . In particular, there exists an $e \in \mathcal{C}_+$ such that $Ke = r(K)e$, and $e \gg 0$.*

Lemma 4.3 follows directly from theorem 1.2 of [Du \(2006\)](#), given that \mathcal{C}_+ is both solid and reproducing. The element e in lemma 4.3 is unique up to a scale factor. In what follows we normalize by requiring that $\|e\| = 1$, and call e the *Perron–Frobenius eigenfunction* of K .

Lemma 4.4. *If K is strongly positive and compact, and ϕ and K jointly satisfy*

$$(20) \quad \lim_{t \downarrow 0} \frac{\phi(t)}{t} r(K) > 1 \quad \text{and} \quad \lim_{t \uparrow \infty} \frac{\phi(t)}{t} r(K) < 1,$$

then there exist positive constants $c_1 < c_2$ with the following properties:

- (a) *If $0 < c \leq c_1$ and $f = ce$, then $f \ll Af$.*
- (b) *If $c_2 \leq c < \infty$ and $f = ce$, then $Af \ll f$.*

Proof. Let $\lambda := r(K)$. Since K is strongly positive and compact, the Perron–Frobenius eigenfunction e discussed above is well defined. Regarding claim (a), observe that, in view of (20), there exists an $\epsilon > 0$ such that

$$\frac{\phi(t)}{t} \lambda > 1 \quad \text{whenever} \quad 0 < t < \epsilon.$$

Choosing c_1 such that $0 < c_1 < \epsilon/\lambda$ and $c \leq c_1$, we have $c\lambda e(x) \leq c_1 \lambda \|e\| = c_1 \lambda < \epsilon$, and hence

$$Ace(x) = \phi(cKe(x)) = \phi(c\lambda e(x)) = \frac{\phi(c\lambda e(x))}{c\lambda e(x)} c\lambda e(x) > ce(x).$$

Since $x \in \mathbb{X}$ was arbitrary, the first claim in the lemma is verified.

Turning to claim (b) and using again the hypotheses in (20), we can choose a finite M such that

$$\frac{\phi(t)}{t} \lambda < 1 \quad \text{whenever} \quad M < t.$$

Let m be the minimum of e on \mathbb{X} . Since \mathbb{X} is compact and $e \gg 0$, we have $m > 0$. Let c_2 be a constant strictly greater than $\max\{M/(\lambda m), c_1\}$ and let c lie in $[c_2, \infty)$.

By the definition of m we have $c\lambda e(x) \geq c_2\lambda e(x) > M$ for all $x \in \mathbb{X}$, from which it follows that

$$Ace(x) = \phi(c\lambda e(x)) = \frac{\phi(c\lambda e(x))}{c\lambda e(x)} \lambda ce(x) < ce(x).$$

By construction, $0 < c_1 < c_2$, so all claims are now established. \square

Proposition 4.2. *Let K be strongly positive and compact, and let the conditions in (20) hold. If, in addition, ϕ is increasing and concave, then A is globally asymptotically stable on \mathcal{C}_+ .*

Proof. Given that ϕ is increasing and concave on \mathbb{R}_+ , lemma 4.1 implies that A is increasing and concave on \mathcal{C}_+ . Since lemma 4.4 implies existence of a pair f_1, f_2 such that $Af_1 \gg f_1$ and $Af_2 \leq f_2$, and since the function f_1 can be chosen from $\mathring{\mathcal{C}}_+$, theorem 4.1 implies that A has a fixed point g^* in $\mathring{\mathcal{C}}_+$.

Next we claim that

$$(21) \quad \forall g \in \mathring{\mathcal{C}}_+, \exists f_1, f_2 \in \mathring{\mathcal{C}}_+ \text{ such that } f_1 \leq g, g^* \leq f_2, Af_1 \gg f_1 \text{ and } Af_2 \leq f_2.$$

To see this, fix $g \in \mathring{\mathcal{C}}_+$. Since $g \gg 0$ and \mathbb{X} is compact, g attains a finite maximum and strictly positive minimum on \mathbb{X} . The same is true of the existing fixed point g^* and the Perron–Frobenius eigenfunction e . Hence, we can choose constants a_1 and a_2 such that $0 \ll a_1 e \leq g^*, g \leq a_2 e$. With a_1 chosen sufficiently small and a_2 sufficiently large, lemma 4.4 implies that $a_1 e \ll A(a_1 e)$ and $A(a_2 e) \ll a_2 e$. Thus (21) holds.

Turning to uniqueness, let g^{**} be a second fixed point of A in $\mathring{\mathcal{C}}_+$. By (21) there exist $f_1, f_2 \in \mathring{\mathcal{C}}_+$ such that $f_1 \leq g^{**}, g^* \leq f_2$ with $f_1 \ll Af_1$ and $Af_2 \ll f_2$. By theorem 4.1, the interval $[f_1, f_2]$ contains only one fixed point. Thus, $g^* = g^{**}$.

Finally, regarding convergence, let g be an element of $\mathring{\mathcal{C}}_+$. Invoking (21) establishes the existence of $f_1, f_2 \gg 0$ such that $f_1 \leq g, g^* \leq f_2$ with $f_1 \ll Af_1$ and $Af_2 \ll f_2$. By theorem 4.1, every element of $[f_1, f_2]$ converges to g^* under iteration of A . In particular, $A^n g \rightarrow g^*$ as $n \rightarrow \infty$. \square

5. PROOFS PART II: THE RECURSIVE UTILITY PROBLEM

5.1. A Topologically Conjugate Problem. Rather than studying T directly, we study a continuous transformation of T and apply the concept of topological conjugacy. In particular, consider the operator A on $\mathring{\mathcal{C}}_+$ defined by

$$(22) \quad Ag(x) = \left\{ \zeta + (Kg(x))^{1/\theta} \right\}^\theta.$$

Let Θ be the transformation defined by $(\Theta w)(x) = w^\theta(x) := w(x)^\theta$. Since $t \mapsto t^\theta$ is a homeomorphism from $(0, \infty)$ to itself, the operator Θ is a homeomorphism from \mathcal{C}_+° to itself.

Lemma 5.1. *T and A are topological conjugate on \mathcal{C}_+° under Θ .*

Proof. We need to show that $\Theta Tw = A\Theta w$ for all $w \in \mathcal{C}_+^\circ$. To see this, fix $w \in \mathcal{C}_+^\circ$ and observe that, for any $x \in \mathbb{X}$,

$$(\Theta Tw)(x) = \left\{ \zeta + (Kw^\theta(x))^{1/\theta} \right\}^\theta = (Aw^\theta)(x) = A\Theta w(x). \quad \square$$

This means that global asymptotic stability of one of these operators on \mathcal{C}_+° is logically equivalent to global asymptotic stability of the other. The next lemma slightly strengthens this result in terms of implications for T .

Lemma 5.2. *If A is globally asymptotically stable on \mathcal{C}_+° , then T is globally asymptotically stable on \mathcal{C}_+ and the fixed point lies in \mathcal{C}_+° .*

Proof. Let A be globally asymptotically stable on \mathcal{C}_+° . From lemma 5.1, it follows that T is globally asymptotically stable on \mathcal{C}_+° . Moreover, since T maps \mathcal{C}_+ into \mathcal{C}_+° , we know that T has no fixed point in $\mathcal{C}_+ \setminus \mathcal{C}_+^\circ$. The fact that T maps \mathcal{C}_+ into \mathcal{C}_+° also implies that $T^n w$ lies in \mathcal{C}_+° for all $n \geq 1$. Given that T is globally asymptotically stable on \mathcal{C}_+° , this trajectory converges to the fixed point $w^* \in \mathcal{C}_+^\circ$. In particular, T is globally asymptotically stable on all of \mathcal{C}_+ . \square

5.2. Proofs of the Main Theorems.

Proof of theorem 2.1. It suffices to show that $(a) \implies (e) \implies (d) \implies (c) \implies (b) \implies (a)$. Of these, $(e) \implies (d)$ is true by definition, and $(d) \implies (c)$ is obvious (just take w to be the fixed point). We now prove the remainder.

$((a) \implies (e))$ Suppose that $r(K) > 1$ and let A be defined by (22). By lemma 5.2, to establish (e) it suffices to show that A is globally asymptotically stable on \mathcal{C}_+° . To see that this is so, let ϕ be the map from \mathbb{R}_+ to itself defined by

$$(23) \quad \phi(t) = \left\{ \zeta + t^{1/\theta} \right\}^\theta$$

with $\phi(0) = 0$. With this definition of ϕ we can express A as $Ag(x) = \phi(Kg(x))$ for $g \in \mathcal{C}_+^\circ$ and $x \in \mathbb{X}$. To complete the proof of (e), we need only show that the conditions of proposition 4.2 hold with ϕ as defined in (23) and K as given in (9).

The operator K is strongly positive by assumption. Under the assumption that $\theta < 0$, the map ϕ is strictly increasing and strictly concave on \mathbb{R}_+ . Thus, we need only show that the two inequalities in (20) are valid. Regarding the first inequality, we have

$$\frac{\phi(t)}{t} = \left\{ \frac{\zeta}{t^{1/\theta}} + 1 \right\}^\theta \uparrow 1 \quad \text{as } t \downarrow 0.$$

Since $r(K) > 1$, the first inequality holds. Regarding the second inequality in (20), evidently $\phi(t)/t \rightarrow 0$ as $t \rightarrow \infty$, so this bound certainly holds. The proof of (e) is therefore complete.

((c) \implies (b)) Let $w \in \mathcal{C}_+$ be such that $\{T^n w\}$ is convergent, with the limit denoted by \bar{w} . Since T maps into the interior of \mathcal{C}_+ , it must be that $\bar{w} \gg 0$. At any such \bar{w} the operator is continuous, being the composition of continuous mappings.¹³ Thus, $T\bar{w} = T(\lim_{n \rightarrow \infty} T^n w) = \lim_{n \rightarrow \infty} T^{n+1} w = \bar{w}$. In particular, \bar{w} is a fixed point of T .

((b) \implies (a)) Suppose to the contrary that $r(K) \leq 1$. It suffices to show that A has no fixed point in $\mathring{\mathcal{C}}_+$. To this end, recall lemma 4.2. We have $r(K) \leq 1$ by assumption and the other conditions (17) have already been checked. Hence lemma 4.2 applies, and A has no fixed point in $\mathring{\mathcal{C}}_+$. \square

Proof of theorem 2.2. As was the case for theorem 2.1, it suffices to show that (a) \implies (e), (c) \implies (b) and (b) \implies (a).

((a) \implies (e)) In view of lemma 5.2, it suffices to show that A given by (22) is globally asymptotically stable on $\mathring{\mathcal{C}}_+$ when $r(K) < 1$. To this end, suppose first that $\theta \in (0, 1)$. The conditions of proposition 4.1 are then satisfied, since $r(K) < 1$ by assumption and ϕ is Lipschitz of order 1. From this proposition we see that A is globally asymptotically stable on \mathcal{C}_+ , with a unique fixed point g^* . This fixed point lies in $\mathring{\mathcal{C}}_+$, since 0 is not a fixed point of A (because $A0 = \phi(K0) = \phi(0) > 0$), and, moreover, if g^* is nonzero then Kg^* is strictly positive, and hence so is $Ag^* = \Phi Kg^*$. Thus, A is also globally asymptotically stable on $\mathring{\mathcal{C}}_+$. Hence (e) is valid.

Now consider the remaining case $\theta \geq 1$, while continuing to assume that $r(K) < 1$. For such θ the function ϕ is increasing and concave, and, since $r(K) < 1$, the conditions in (20) are both satisfied. Hence proposition 4.2 applies, and A is globally asymptotically stable on $\mathring{\mathcal{C}}_+$. Hence (e) holds

¹³Here we are using the fact that K is a compact operator and hence continuous.

((c) \implies (b)) The proof of this implication is identical to the proof of the same implication given in theorem 2.1.

((b) \implies (a)) First we make some observations about $\phi(t) = (\zeta + t^{1/\theta})^\theta$ on \mathbb{R}_+ when $\theta > 0$. Evidently ϕ is continuous and increasing with $\phi(t) > t$ for all $t \in \mathbb{R}_+$. It is not difficult to see that, in addition,

$$(24) \quad 0 \leq s \leq t \implies \phi(s) \leq \frac{\phi(t)}{t}s \quad \text{and} \quad \lim_{n \rightarrow \infty} \phi^n(t) = \infty, \quad \forall t \geq 0.$$

Now suppose that T has a fixed point $g \in \mathcal{C}_+$ and yet $r(K) \geq 1$. Observe that g must lie in $\mathring{\mathcal{C}}_+$, since T maps \mathcal{C}_+ into $\mathring{\mathcal{C}}_+$. Let e be the Perron–Frobenius eigenfunction of K and let c be a positive constant such that $e_c := ce \leq g$. Such a c exists because $\min_{x \in \mathbb{X}} g(x)$ is strictly positive and $\max_{x \in \mathbb{X}} e(x)$ is finite. Let t_0 be a positive constant such that $e_c \leq t_0$ on \mathbb{X} . We claim that

$$(25) \quad \forall n \in \mathbb{N}, \quad A^n e_c \geq \frac{\phi^n(t_0)}{t_0} e_c \quad \text{on } \mathbb{X}.$$

To see this, observe that (25) holds at $n = 0$. Now suppose that (25) holds at some $n \geq 0$. We then have

$$A^{n+1} e_c(x) = \phi(KA^n e_c(x)) \geq \phi\left(\frac{\phi^n(t_0)}{t_0} K e_c(x)\right) \geq \phi\left(\frac{\phi^n(t_0)}{t_0} e_c(x)\right).$$

Here the first inequality is by the induction hypothesis, the monotonicity of ϕ and K and the linearity of K . The second is from $r(K) \geq 1$, which gives $K e_c = c K e = cr(K)e \geq ce = e_c$. Using $e_c(x) \leq t_0$ and the first property in (24), we have

$$A^{n+1} e_c(x) \geq \phi\left(\frac{\phi^n(t_0)}{t_0} e_c(x)\right) \geq \frac{\phi(\phi^n(t_0))}{\phi^n(t_0)} \frac{\phi^n(t_0)}{t_0} e_c(x) = \frac{\phi^{n+1}(t_0)}{t_0} e_c(x).$$

Thus, the statement in (25) is valid.

From (25) and the second property in (24) we conclude that $A^n e_c$ diverges to $+\infty$. Moreover, the fixed point g satisfies $g \geq e_c$, so $A^n g \geq A^n e_c$. Hence $A^n g$ eventually exceeds g , contradicting our assumption that g is a fixed point of A . \square

REFERENCES

- Aliprantis, C. D. and Border, K. C. (2006). *Infinite dimensional analysis*. Springer, 3rd edition.
- Alvarez, F. and Jermann, U. J. (2005). Using asset prices to measure the persistence of the marginal utility of wealth. *Econometrica*, 73(6):1977–2016.

- Balbus, L. (2016). On non-negative recursive utilities in dynamic programming with nonlinear aggregator and CES. Working paper, Available at SSRN: <https://ssrn.com/abstract=2703975>.
- Bansal, R., Kiku, D., Yaron, A., et al. (2012). An empirical evaluation of the long-run risks model for asset prices. *Critical Finance Review*, 1(1):183–221.
- Bansal, R. and Yaron, A. (2004). Risks for the long run: A potential resolution of asset pricing puzzles. *The Journal of Finance*, 59(4):1481–1509.
- Becker, R. A., Boyd, J. H., and Sung, B. Y. (1989). Recursive utility and optimal capital accumulation. i. existence. *Journal of Economic Theory*, 47(1):76–100.
- Becker, R. A. and Rincón-Zapatero, J. P. (2017). Recursive utility and Thompson aggregators. Technical report, University of Indiana Working Paper.
- Bloise, G. and Vailakis, Y. (2017). Convex dynamic programming with (bounded) recursive utility. Technical report, University of Glasgow.
- Boyd, J. H. (1990). Recursive utility and the ramsey problem. *Journal of Economic Theory*, 50(2):326–345.
- Christensen, T. M. (2017). Identification and estimation of dynamic models with robust decision makers. Technical report, New York University.
- Coleman, W. J. (1991). Equilibrium in a production economy with an income tax. *Econometrica*, pages 1091–1104.
- Datta, M., Mirman, L. J., and Reffett, K. L. (2002). Existence and uniqueness of equilibrium in distorted dynamic economies with capital and labor. *Journal of Economic Theory*, 103(2):377–410.
- Du, Y. (2006). *Order Structure and Topological Methods in Nonlinear Partial Differential Equations: Maximum principles and applications*. World Scientific.
- Epstein, L. G. and Zin, S. E. (1989). Risk aversion and the temporal behavior of consumption and asset returns: A theoretical framework. *Econometrica*, 57(4):937–969.
- Hansen, L. P., Heaton, J. C., and Li, N. (2008). Consumption strikes back? measuring long-run risk. *Journal of Political Economy*, 116(2):260–302.
- Hansen, L. P. and Scheinkman, J. A. (2009). Long-term risk: An operator approach. *Econometrica*, 77(1):177–234.
- Hansen, L. P. and Scheinkman, J. A. (2012). Recursive utility in a markov environment with stochastic growth. *Proceedings of the National Academy of Sciences*, 109(30):11967–11972.
- Kolmogorov, A. and Fomin, S. (1975). Introductory real analysis.
- Koopmans, T. C. (1960). Stationary ordinal utility and impatience. *Econometrica*, pages 287–309.

- Krasnosel'skii, M., Vainikko, G., Zabreyko, R., Ruticki, Y., and Stet'senko, V. (2012). *Approximate Solution of Operator Equations*. Springer Netherlands.
- Kreps, D. M. and Porteus, E. L. (1978). Temporal resolution of uncertainty and dynamic choice theory. *Econometrica*, pages 185–200.
- Le Van, C. and Vailakis, Y. (2005). Recursive utility and optimal growth with bounded or unbounded returns. *Journal of Economic Theory*, 123(2):187–209.
- Lucas, R. E. and Stokey, N. L. (1984). Optimal growth with many consumers. *Journal of Economic Theory*, 32(1):139–171.
- Marinacci, M. and Montrucchio, L. (2010). Unique solutions for stochastic recursive utilities. *Journal of Economic Theory*, 145(5):1776–1804.
- Mirman, L. J., Morand, O. F., and Reffett, K. L. (2008). A qualitative approach to markovian equilibrium in infinite horizon economies with capital. *Journal of Economic Theory*, 139(1):75–98.
- Ozaki, H. and Streufert, P. A. (1996). Dynamic programming for non-additive stochastic objectives. *Journal of Mathematical Economics*, 25(4):391–442.
- Qin, L. and Linetsky, V. (2017). Long-term risk: A martingale approach. *Econometrica*, 85(1):299–312.
- Rincón-Zapatero, J. P. and Rodríguez-Palmero, C. (2007). Recursive utility with unbounded aggregators. *Economic Theory*, 33(2):381–391.
- Schorfheide, F., Song, D., and Yaron, A. (2017). Identifying long-run risks: A bayesian mixed-frequency approach. Technical report, University of Pennsylvania working paper.
- Streufert, P. A. (1990). Stationary recursive utility and dynamic programming under the assumption of biconvergence. *The Review of Economic Studies*, 57(1):79–97.
- Wagner, C. H. (1982). A generic approach to iterative methods. *Mathematics Magazine*, 55(5):259–273.
- Weil, P. (1990). Nonexpected utility in macroeconomics. *The Quarterly Journal of Economics*, 105(1):29–42.
- Zaanen, A. C. (1997). *Introduction to Operator Theory in Riesz Spaces*. Springer.
- Zhang, Z. (2013). *Variational, topological, and partial order methods with their applications*. Springer.